

VAC-SULES REPRODUCTIVA H

Double emulsion combined vaccine



Composition:

Infectious Bovine Rhinotracheitis virus type 1 (IBR) $\geq 1 \times 10^7$ DICT50 % / mL
Infectious Bovine Rhinotracheitis virus type 5 (IBR) $\geq 1 \times 10^7$ DICT50 % / mL
Bovine Viral Diarrhea virus type 1 (BVD) $\geq 1 \times 10^5$ DICT50 % / mL
Bovine Viral Diarrhea virus type 2 (BVD) $\geq 1 \times 10^5$ DICT50 % / mL
Leptospira Icterohaemorrhagiae $\geq 1 \times 10^7$ UFC/ mL
Leptospira Canicola $\geq 1 \times 10^7$ UFC/ mL
Leptospira Copenhageni $\geq 1 \times 10^7$ UFC/ mL
Leptospira Pomona $\geq 1 \times 10^7$ UFC/ mL
Leptospira Wolffi $\geq 1 \times 10^7$ UFC/ mL
Leptospira Grippotyphosa $\geq 1 \times 10^7$ UFC/ mL
Leptospira Tarassovi $\geq 1 \times 10^7$ UFC/ mL
Leptospira Hardjo $\geq 1 \times 10^7$ UFC/ mL
Leptospira Pyrogenes $\geq 1 \times 10^7$ UFC/ mL
Leptospira Bataviae $\geq 1 \times 10^7$ UFC/ mL
Leptospira Hardjo bovis $\geq 1 \times 10^7$ UFC/ mL
Campylobacter fetus subespecie fetus 40 mg / dose
Campylobacter fetus subespecie venerealis 40 mg / dose
Campylobacter fetus subespecie venerealis biotipo intermedius 40 mg / dose
Haemophilus somnus $\geq 1 \times 10^7$ UFC/ mL

Excipients:

Aluminum hydroxyde 10 % v/v
Saline solution (NaCl 0.85 % w/v) q.s. dose
Thimerosal 0.5 mg / dose

Indications for use:

Vaccine for the prevention of Infectious Bovine Rhinotracheitis, Bovine Viral Diarrhea, Leptospirosis, Campylobacteriosis and Haemofilosis in bovine.

Target species:

Bovine

Withdrawal period:

Do not slaughter animals for human consumption before 45 – 60 days post-vaccination.

Route of administration:

Subcutaneous route.

Dosage:

Bovine: 5 mL
Vaccinate and revaccinated at an interval of 20-25 days, and revaccinate with a single dose annually.
It can be vaccinated animals from three months old. It can be used in pregnant females.

Conservation: 2°C – 8°C. DO NOT FREEZE.