

VAC-SULES LEPTO 10 H

Aluminium hydroxide bacterial vaccine

Composition:

Leptospira Icterohaemorrhagiae $\geq 1 \times 10^7$ UFC/ mL

Leptospira Canicola $\geq 1 \times 10^7$ UFC/ mL

Leptospira Copenhageni $\geq 1 \times 10^7$ UFC/ mL

Leptospira Pomona $\geq 1 \times 10^7$ UFC/ mL

Leptospira Wolffi $\geq 1 \times 10^7$ UFC/ mL

Leptospira Grippotyphosa $\geq 1 \times 10^7$ UFC/ mL

Leptospira Tarassovi $\geq 1 \times 10^7$ UFC/ mL

Leptospira Hardjo $\geq 1 \times 10^7$ UFC/ mL

Leptospira Pyrogenes $\geq 1 \times 10^7$ UFC/ mL

Leptospira Bataviae $\geq 1 \times 10^7$ UFC/ mL



Excipients:

Aluminium hydroxide 10 % v/v

Saline Solution (NaCl 0.85 % w/v) q.s. dose

Thimerosal 0.5 mg / dose

Indications for use:

Vaccine for the prevention of Leptospirosis in bovine, ovine, caprine and swine.

Target species:

Bovine, ovine, caprine and swine.

Withdrawal period:

Do not slaughter animals for human consumption before 45 -60 days post-vaccination.

Route of administration:

Subcutaneous route.

Dosage:

Bovine, ovine, caprine and swine: 2 mL

Vaccinate healthy and stress-free animals with two doses with an interval of 20-25 days, and

revaccinate with a single dose annually.

In cases of high incidence of the disease, vaccinate twice a year or according to the veterinarian's discretion.

It can be applied to pregnant females.

Conservation:

4 °C – 8°C. DO NOT FREEZE.