

VAC-SULES LEPTO 11

Double emulsion bacterial vaccine

Composition:

Leptospira Icterohaemorrhagiae $\geq 1 \times 10^7$ UFC/ mL

Leptospira Canicola $\geq 1 \times 10^7$ UFC/ mL

Leptospira Copenhageni $\geq 1 \times 10^7$ UFC/ mL

Leptospira Pomona $\geq 1 \times 10^7$ UFC/ mL

Leptospira Wolffi $\geq 1 \times 10^7$ UFC/ mL

Leptospira Grippotyphosa $\geq 1 \times 10^7$ UFC/ mL

Leptospira Tarassovi $\geq 1 \times 10^7$ UFC/ mL

Leptospira Hardjo $\geq 1 \times 10^7$ UFC/ mL

Leptospira Pyrogenes $\geq 1 \times 10^7$ UFC/ mL

Leptospira Bataviae $\geq 1 \times 10^7$ UFC/ mL

Leptospira Hardjo bovis $\geq 1 \times 10^7$ UFC/ mL

Excipients:

Double emulsion adjuvant 0.75 mL / dose

Thimerosal 1/10,000



Indications for use:

Vaccine for the prevention of Leptospirosis in bovine, ovine, caprine and swine.

Target species:

Bovine, ovine, caprine and swine.

Withdrawal period:

Do not slaughter animals for human consumption before 45 -60 days post-vaccination.

Route of administration:

Subcutaneous route.

Dosage:

Bovine: 3 mL. Ovine, caprine and swine: 2 mL

Vaccinate healthy and stress-free animals with two doses with an interval of 20-25 days, and revaccinate with a single dose annually. It can be vaccinated animals from three months old. In cases of high incidence of the disease, vaccinate twice a year or according to the veterinarian's discretion.

It can be applied to pregnant females.

Conservation: 2°C – 8°C. DO NOT FREEZE.