

VAC-SULES POLIVAC 10/1

Clostridial vaccine

Composition:

Purified toxoids of:

Clostridium novyi type B ≥ 500 DLM /mL

Clostridium perfringens type D ≥ 200 L+ /mL

Clostridium perfringens type B ≥ 200 L+ /mL

Clostridium perfringens type C ≥ 200 L+ /mL

Clostridium perfringens type A ≥ 20 DLM/mL

Clostridium septicum ≥ 100 DLM/mL

Clostridium sordellii ≥ 500 DLM/mL

Clostridium tetani $\geq 300,000$ DLM/mL

Integral anaerobes of:

Clostridium chauvoei ≥ 8 μ L packed cells

Clostridium haemolyticum ≥ 0.4 UH/mL

Excipients:

Aluminium Hydroxide 10 % v/v

Saline solution (NaCl 0,85 % p/v) q.s.dose

Thimerosal 0.5 mg / dose



Indications for use:

Vaccine for the prevention of Hemorrhagic Gastritis, Gas Gangrene, Lambs Dysentery, Sheep Enterotoxaemia, Pulpy Kidney Disease, Necrotizing Infectious Hepatitis, Bacillary Hemoglobinuria, Sudden Death, Tetanus and Symptomatic Anthrax.

Target species:

Bovine, ovine and caprine

Withdrawal period:

Do not slaughter animals for human consumption before 25 days post-vaccination.

Route of administration:

Subcutaneous route.

Dosage:

Bovine: 5 mL

Ovine and caprine: 3 mL

Vaccinate and revaccinated at an interval of 28-30 days, and revaccinate with a single dose annually.

It can be vaccinated animals from two months old.

Conservation:

2°C – 8°C. DO NOT FREEZE.